

# Medicine Labels for “A Case of Unintentional Overdose”

## Drug Facts

<b>Active Ingredients</b> (in each caplet) Acetaminophen 500 mg.....	<b>Purpose</b> Pain reliever/ fever reducer
<b>Uses:</b> temporarily relieves minor aches and pains due to:	
<ul style="list-style-type: none"> <li>• headaches</li> <li>• backache</li> <li>• menstrual cramps</li> </ul>	<ul style="list-style-type: none"> <li>• muscle aches</li> <li>• arthritis</li> <li>• toothache</li> <li>• the common cold</li> </ul>
<b>Warnings</b>	
<b>Alcohol warning:</b> If you consume 3 or more alcoholic beverages every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.	
<b>Do not use</b> with any other products containing acetaminophen	
<b>Stop use and ask your doctor if</b>	
<ul style="list-style-type: none"> <li>• new symptoms occur</li> <li>• redness or swelling is present</li> <li>• pain gets worse or lasts for more than 10 days</li> <li>• fever gets worse or lasts for more than three days</li> </ul>	
If pregnant or breast-feeding, ask a health professional before use.	
<b>Keep out of reach of children.</b>	

*Adult*

**Extra Strength**

**Pain Reliever  
and Fever Reducer**

*Contains*

**Acetaminophen**

**Caplets** 

**100 CAPLETS—500 mg each**

<b>Overdose warning:</b> Taking more than the recommended dose (overdose) could cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
<b>Directions.</b>
<ul style="list-style-type: none"> <li>• do not take more than directed (see overdose warning)</li> <li>• take 2 caplets every 4 to 6 hours as needed</li> <li>• do not take more than 8 caplets in 24 hours</li> </ul>
<b>Children under 12 years:</b> do not use this adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and could cause serious health problems.
<b>Other Information</b>
<ul style="list-style-type: none"> <li>• do not use if neck wrap or foil inner seal is broken or missing</li> <li>• store at room temperature</li> </ul>
<b>Questions or comments?</b> Call toll free 1-800-XXX-XXXX

## Drug Facts

### Active Ingredients

(in 15 ml TBSP)

### Purposes

Acetaminophen, 325 mg..... Fever reducer/pain reliever  
Dextromethorphan HBr, 15 mg..... Cough suppressant  
Doxylamine Succinate, 625 mg.....Antihistamine

### Uses:

- temporarily relieves common cold/flu symptoms
- cough due to minor throat and bronchial irritation
- throat
- headache
- fever
- minor aches and pains
- runny nose and sneezing

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in any 24-hour period, which is the maximum daily amount
- other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

### Do not use

- to make a child sleep
- with any prescription or nonprescription drug containing acetaminophen
- if you are taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping these drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this drug.

### Ask a doctor before use if you have:

- liver disease
- glaucoma
- sodium restricted diet
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

### Ask a doctor or pharmacist before use if you are taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

# Nighttime Relief

Helps you sleep when you have a cold or the flu

- Aches, Fever, Sore Throat
- Sneezing, Runny Nose
- Cough

10 fluid ounces (295 ml)  
10% alcohol

### When using this product:

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol and sedatives may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### Stop use and ask a doctor if

- pain or cough gets worse or lasts for more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.

### Keep out of reach of children

**Overdose Warning** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions.

- take only as directed
- use dose cup or tablespoon (TBSP)
- do not exceed 4 doses per 24 hours

•12 years and over	30 ml (2TBSP) every 6 hours
•children 4 to under 12 years	ask a doctor
•children under 4 years	do not use

### Other Information

- each tablespoon contains potassium 5 mg, sodium 18 mg
- store at room temperature

**Inactive ingredients** acesulfame potassium, alcohol, citric acid, D&C yellow no. 10, FD&C green no. 3, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, and sodium citrate.

### Questions or comments?

Call toll free 1-800-XXX-XXXX

# Drug Facts

## Active Ingredients (in each liquid-filled capsule) Purposes

Acetaminophen, 325 mg..... fever reducer/pain reliever  
 Dextromethorphan HBr, 15 mg..... cough suppressant  
 Doxylamine Succinate, 625 mg..... antihistamine

**Uses:** temporarily relieves these symptoms occurring with a cold, flu, hay fever, or other upper respiratory allergies:

- headaches
- sore throat
- itchy, watery eyes
- itchy nose or throat
- minor aches and pains
- cough
- runny nose

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 capsules in any 24-hour period, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

## Do not use

•if you are taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping these drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this drug.  
 •with any prescription or nonprescription drug containing acetaminophen

## Ask a doctor before use if you have:

- liver disease
- glaucoma
- trouble urinating due to enlarged prostate gland
- coughing that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

## Ask a doctor or pharmacist before use if you are taking

- the blood thinning drug warfarin
- any other pain reliever/fever reducer
- sedatives or tranquilizers

# Adult Cold and Flu Formula



## 20 Liquid-Filled Capsules

### Relieves:

- Cough
- Sore Throat
- Fever
- Sneezing
- Body Aches
- Runny Nose

## When using this product:

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol and sedatives may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

## Stop use and ask a doctor if

- pain or cough gets worse or lasts for more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.

## Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions.

- Do not take more than 8 capsules in any 24 hour period
- Do not exceed recommended dosage (Taking more than the recommended dose can cause serious liver damage.

age	dose
•adults and children 12 years and over	2 capsules every 6 hours
•children under 12 years	do not use

## Other Information

- do not use if neck wrap or foil inner seal is broken or missing
- store at room temperature

**Inactive ingredients** FD&C blue no. 1, gelatin, glycerin, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution

## Questions or comments?

Call toll free 1-800-XXX-XXXX